



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

BRAKER2: Incorporating Protein Homology Information into Gene Prediction with GeneMark-EP and AUGUSTUS

A pipeline for fully automated training and prediction

Plant and Animal Genomes XXVI, January 14th 2018

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky

Presenting author: katharina.hoff@uni-greifswald.de

**BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS**

**Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky**



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

Contents

- 1 Gene prediction
- 2 BRAKER1: RNAseq
- 3 BRAKER2: proteins
 - Short evolutionary distance
 - Long evolutionary distance
- 4 Summary
- 5 References

Structural genome annotation problem

Input

- genome assembly
- extrinsic evidence, e.g. from RNAseq, protein database

Output

- protein-coding genes: exon-intron structures (.gff)

Example (from Chr I in *C. elegans*)





BRAKER1: RNAseq integration

BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS FREE

Katharina J. Hoff ✉, Simone Lange, Alexandre Lomsadze, Mark Borodovsky ✉, Mario Stanke

Bioinformatics, Volume 32, Issue 5, 1 March 2016, Pages 767–769,

<https://doi.org/10.1093/bioinformatics/btv661>

- >4000 downloads
- 73 citations since 2016 (google scholar)

BRAKER1: RNAseq integration

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction


BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance
Long evolutionary distance

Summary

References



BRAKER2: Part I - proteins of closely related species

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



[Gene prediction](#)

[BRAKER1: RNAseq](#)


[BRAKER2: proteins](#)

[Short evolutionary distance](#)

[Long evolutionary distance](#)

[Summary](#)

[References](#)






Drosophila melanogaster and relatives

For a given species,

- the average number of mutations per genomic site was computed from alignments of ortholog gene sequences (including introns).
- the protein identity was computed as average of identity values of the best *exonerate* hit found for each protein of this species against the *D. melanogaster* genome.



Increasing evolutionary distance leads to decreasing gene prediction accuracy of AUGUSTUS

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



[Gene prediction](#)

[BRAKER1: RNAseq](#)

[BRAKER2: proteins](#)


[Short evolutionary distance](#)

[Long evolutionary distance](#)

[Summary](#)

[References](#)

AUGUSTUS ab initio prediction





[Gene prediction](#)

[BRAKER1: RNAseq](#)

[BRAKER2: proteins](#)

[Short evolutionary distance](#)


[Long evolutionary distance](#)

[Summary](#)

[References](#)

Increasing evolutionary distance leads to decreasing gene prediction accuracy of AUGUSTUS

AUGUSTUS prediction with training set hints





Increasing evolutionary distance leads to decreasing gene prediction accuracy of AUGUSTUS

With increasing distance between query protein and target genome, spliced alignments become

- less sensitive while keeping a constant level of specificity (e.g. GenomeThreader),
- or both less sensitive and less specific (e.g. Exonerate).

Therefore, training AUGUSTUS on spliced alignments is suitable upon availability of a very closely related query species, only!

BRAKER2: Part II - proteins of more remote species

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins


Short evolutionary distance

Long evolutionary distance

Summary

References

“Standard mapping approach”: proteins to genome



→ works well for closely related species, only

BRAKER2: Part II - proteins of more remote species

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq


BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References



Protein database for gene prediction in *D. melanogaster*

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

Insect portion of EggNOG (inNOG) excluding *Drosophila* species

- *Acyrtosiphon pisum*
- *Aedes aegypti*
- *Anopheles darlingi*
- *Anopheles gambiae*
- *Apis mellifera*
- *Atta cephalotes*
- *Bombyx mori*
- *Culex quinquefasciatus*
- *Danaus plexippus*
- *Heliconius melpomene*
- *Nasonia vitripennis*
- *Pediculus humanus*
- *Tribolium castaneum*



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance


Summary

References

Intron recovery from protein mapping

Protein mapping with no *Drosophila* EggNOG (inNOG)

- 30,996 introns predicted
- 21,843 matched introns in CDS part of the annotated genes



Mapping of proteins from remote species recovers $\sim 45\%$ of introns with specificity of $\sim 70\%$.



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary


References

Intron recovery from protein mapping

Protein mapping with some *Drosophila* species present as external evidence

no_Dro
w_gvw
w_gvwpa

no *Drosophila* species
with *D. grimshawi*, *D. virilis*, *D. willistoni*
with *D. grimshawi*, *D. virilis*, *D. willistoni*, *D. pseudoobscura*, *D. ananassae*



- more introns were detected
- performance of protein mapping with addition of 5 fly proteomes came closer to performance with RNAseq external evidence

Accuracy of GeneMark-EX with different sources of evidence

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins


Short evolutionary distance

Long evolutionary distance

Summary

References



- results are on **softmasked genome** (strongly recommended!)



- GeneMark-EP and GeneMark-ET outperformed GeneMark-ES
- GeneMark-EP with “remote” proteins was comparable with GeneMark-ET
- GeneMark-EP and GeneMark-ET were close to the best possible performance: compared to training with “ideal” introns

Accuracy of BRAKER2

Gene prediction accuracy (F1)



**BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS**

**Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky**



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

Summary

- BRAKER2 is a novel fully automatic pipeline which makes gene prediction in eukaryotic genomes with RNAseq or protein external evidence.
- Training in BRAKER2 is done by GeneMark-EX which particularly can use remote proteins as external evidence.
- Prediction in BRAKER2 is done by AUGUSTUS using RNAseq or proteins as hints.

**BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS**

**Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky**



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

Ongoing & future work

- Optimization of evidence integration in BRAKER2
- Combining RNAseq and protein information
- UTR training & integration of RNAseq coverage information



References


- Hoff, Katharina J., et al. "BRAKER1: unsupervised RNAseq-based genome annotation with GeneMark-ET and AUGUSTUS." *Bioinformatics* 32.5 (2015): 767-769.
- Stanke, Mario, et al. "Using native and syntenically mapped cDNA alignments to improve de novo gene finding." *Bioinformatics* 24.5 (2008): 637-644.
- Lomsadze, Alexandre, Paul D. Burns, and Mark Borodovsky. "Integration of mapped RNAseq reads into automatic training of eukaryotic gene finding algorithm." *Nucleic acids research* 42.15 (2014): e119-e119.
- Slater, Guy St C., and Ewan Birney. "Automated generation of heuristics for biological sequence comparison." *BMC bioinformatics* 6.1 (2005): 31.
- Gremme, Gordon. "GenomeThreader Gene Prediction Software." (2014).
- Dobin, Alexander, et al. "STAR: ultrafast universal RNA-seq aligner." *Bioinformatics* 29.1 (2013): 15-21.

BRAKER2 is available for download at

- <http://bioinf.uni-greifswald.de>
- <http://exon.gatech.edu>

State of the art: BRAKER with RNAseq & proteins

Close homology



BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance
Long evolutionary distance

Summary

References

State of the art: BRAKER with RNAseq & proteins

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



[Gene prediction](#)

[BRAKER1: RNAseq](#)

[BRAKER2: proteins](#)


Short evolutionary distance

Long evolutionary distance

[Summary](#)

[References](#)

AUGUSTUS ab initio prediction



State of the art: BRAKER with RNAseq & proteins

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



[Gene prediction](#)

[BRAKER1: RNAseq](#)

[BRAKER2: proteins](#)


Short evolutionary distance

Long evolutionary distance

[Summary](#)


[References](#)

AUGUSTUS prediction with training set hints



State of the art: BRAKER with RNAseq & proteins

Remote homology



BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



Gene prediction

BRAKER1: RNAseq

BRAKER2: proteins

Short evolutionary distance

Long evolutionary distance

Summary

References

State of the art: BRAKER with RNAseq & proteins

BRAKER2:
Incorporating
Protein Homology
Information into
Gene Prediction with
GeneMark-EP and
AUGUSTUS

Katharina J. Hoff,
Alexandre Lomsadze,
Mario Stanke,
Mark Borodovsky



[Gene prediction](#)

[BRAKER1: RNAseq](#)

[BRAKER2: proteins](#)

Short evolutionary distance

Long evolutionary distance

[Summary](#)

[References](#)

